arXiv:1611.03667v2 [math.AC] 14 Nov 2016

Ring of Real Analytic Functions on $[0, 1]$

Sagar Shrivastava & Vaibhav Pandey

Abstract

We consider the ring of real analytic functions defined on $[0, 1]$, i.e.

$$C^{\omega}[0, 1] = \{f : [0, 1] \longrightarrow \mathbb{R} | f \text{ is analytic on } [0, 1]\}$$

In this article, we explore the nature of ideals in this ring. It is well known that the ring $C[0, 1]$ of real valued continuous functions on $[0, 1]$ has precisely the following maximal ideals:

$$\text{For } \gamma \in [0, 1], M_{\gamma} := \{f \in C[0, 1] | f(\gamma) = 0\}$$

It has been proved that each such M_{γ} is infinitely generated, in-fact uncountably generated [1]. Observe that $C^{\omega}[0, 1]$ is a subring of $C[0, 1]$

We prove that for any γ in $[0, 1]$, the contraction M_{γ}^{ω} of M_{γ} under the natural inclusion of $C^{\omega}[0, 1]$ in $C[0, 1]$ is again a maximal ideal (of $C^{\omega}[0, 1]$), and these are precisely all the maximal ideals of $C^{\omega}[0, 1]$. Next we prove that each M_{γ}^{ω} is principal (though M_{γ} is uncountably generated). Surprisingly, this forces all the ideals of the ring $C^{\omega}[0, 1]$ to be singly generated, i.e. $C^{\omega}[0, 1]$ is a PID.

To begin with, we observe that the maximal ideals of $C^\omega[0,1]$ are precisely $M_\gamma^\omega(\gamma \in [0,1])$.

Firstly, M_γ^ω is the kernel of the following surjective homomorphism:

$$\begin{aligned} \Phi_\gamma : C^\omega[0,1] &\longrightarrow \mathbb{R} \\ f &\mapsto f(\gamma) \end{aligned}$$

Hence $C^\omega[0,1]/M_\gamma^\omega \cong \mathbb{R}$. Therefore M_γ^ω is maximal.

Further, the fact that any maximal ideal of $C^\omega[0,1]$ is of the form M_γ^ω for some $\gamma \in [0,1]$ is merely a consequence of compactness of $[0,1]$. The proof for this is precisely the proof of the fact that any maximal ideal of $C[0,1]$ is of the form M_γ.

Theorem 1. *M_γ^ω is a principal ideal generated by $(x-\gamma)$*

Proof. Before we come to the main proof, note that

$$M_\gamma^\omega = \{f(x) - f(\gamma) | f \in C^\omega[0,1]\}$$

Let f be any function in $C^\omega[0,1]$, we define

$$g(x) := \frac{f(x) - f(\gamma)}{x - \gamma}$$

We claim that g is also analytic on $[0,1]$. For a moment, assume the claim. Then we have $f(x) - f(\gamma) = (x-\gamma)g(x)$, and hence any arbitrary element f of M_γ^ω is contained in the ideal generated by $(x-\gamma)$, hence M_γ^ω is a principal ideal.

So now let us prove the claim. We prove our claim in two parts. First we show that there exists a convergent Taylor series for g in a neighborhood of γ and then we show that in any neighborhood not containing γ, it is analytic, and hence analytic on the whole of $[0,1]$.

Since f is analytic, we can get a Taylor series expansion of f at γ. Then for x in a neighborhood of γ, we have-

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} f^{(n)}(\gamma) \frac{(x-\gamma)^n}{n!} \\ \Rightarrow f(x) - f(\gamma) &= (x-\gamma) \sum_{n=1}^{\infty} f^{(n)}(\gamma) \frac{(x-\gamma)^{n-1}}{n!} \\ \Rightarrow \frac{f(x) - f(\gamma)}{x-\gamma} &= \sum_{n=1}^{\infty} f^{(n)}(\gamma) \frac{(x-\gamma)^{n-1}}{n!} = \sum_{n=0}^{\infty} \frac{f^{(n+1)}(\gamma)}{n+1} \frac{(x-\gamma)^n}{n!} \end{aligned}$$

The LHS of the above equation is $g(x)$ by definition, and we can say that the RHS is a Taylor series expansion of $g(x)$ where $g^{(n)}(\gamma) = \frac{f^{(n+1)}(\gamma)}{n+1}$. We claim that the RHS is convergent in a neighborhood of γ. Since f is analytic, its derivative (f') is also analytic on a neighborhood around γ, and we get the following Taylor series expansion at γ given by

$$f'(x) = \sum_{n=0}^{\infty} f^{(n+1)}(\gamma)\frac{(x-\gamma)^n}{n!}$$

Clearly, the coefficients of $\frac{(x-\gamma)^n}{n!}$ in the expansion of f' and g satisfy the following inequality in the intersection of the neighborhoods of γ where f and f' are expressed in their Taylor series-

$$|g^{(n)}(\gamma)| = \frac{|f^{(n+1)}(\gamma)|}{n+1} \leq |f^{(n+1)}(\gamma)|$$

Since the Taylor series of f' converges in this neighborhood, it forces the Taylor series of g to converge in this neighborhood of γ. Hence g is analytic in some neighborhood of γ (call that neighborhood N, where $N = [0, \varepsilon)$ if $\gamma = 0$, $(1-\delta, 1]$ if $\gamma = 1$, $(\gamma - \delta, \gamma + \varepsilon)$ otherwise, for $\varepsilon, \delta > 0$)

For the other part of the claim, we show that g is analytic on the set N' ($N' = (\varepsilon/2, 1]$ if $\gamma = 0$, $N' = [0, 1-\delta/2)$ if $\gamma = 1$, $N' = [0, \gamma-\delta/2) \cup (\gamma+\varepsilon/2, 1]$ otherwise, where $\varepsilon, \delta > 0$ are the same as used for N)

Now $(x - \gamma)$ is analytic on N', and $x - \gamma \neq 0$ in N'.
$\therefore \frac{1}{(x-\gamma)}$ is analytic on N', and $f(x) - f(\gamma)$ is also analytic on N' (as it is analytic on $[0, 1]$). So,

$$g(x) = (f(x) - f(\gamma))\frac{1}{(x-\gamma)} = \frac{f(x) - f(\gamma)}{x - \gamma}$$

As g is a product of two analytic functions on N', it is also an analytic function on N'.

Hence g is an analytic function on $N \cup N' = [0, 1]$ So we get that M_γ^ω is a principal ideal. □

Theorem 2. *$C^\omega[0, 1]$ is an integral domain.*

Proof. Let $f, g \in C^\omega[0, 1]$ be such that $f.g \equiv 0$. We will use the fact that an analytic function is if and only if all the coefficients in the Taylor series expansion are zero. So we expand the functions f, g along $\frac{1}{2}$, without loss of generality assume that $f \not\equiv 0$, hence we will show that $g \equiv 0$. But this will be the same as proving that power series ring over an integral domain is again an integral domain, which proves that $C^\omega[0, 1]$ is an integral domain. □

Corollary. *$C^\omega[0,1]$ is a PID.*

Proof. Let I be a non-zero proper ideal of $C^\omega[0,1]$. Then it must be contained in some maximal ideal. Then by (1), $I \subseteq \langle x-\gamma_1\rangle$ for some $\gamma_1 \in [0,1]$.

Hence $I = \langle x-\gamma_1\rangle I'$, for some ideal I' of $C^\omega[0,1]$. If $I' = C^\omega[0,1]$, $I = \langle x-\gamma_1\rangle = M^\omega_{\gamma_1}$ and we are done.

If not, then I' is a proper ideal of $C^\omega[0,1]$ and $I' \subseteq \langle x-\gamma_2\rangle$, so that $I = \langle x-\gamma_1\rangle\langle x-\gamma_2\rangle I''$ for some ideal I'' and we continue this process.

We claim that this process will stop after finitely many steps. If not, then choose any non zero function $f \in I$, then $f(\gamma_i) = 0\ \forall i \in \mathbb{N}$. The set $\{\gamma_i\}_{i\in\mathbb{N}}$ is an infinite set in the compact set $[0,1]$, hence by Bolzano-Weistrass theorem, this set has a limit point in $[0,1]$. Also if the zero set of an analytic function has a limit point then the function itself is zero in the connected component containing the limit point. Therefore f is zero in $[0,1]$, which contradicts our assumption that f is nonzero. Hence the process stops after finitely many steps.

Note that the set $\{\gamma_i\}_{i\in\mathbb{N}}$ may have repetitions, but it does not affect our argument.

Hence $I = \langle x-\gamma_1\rangle\langle x-\gamma_2\rangle \dots \langle x-\gamma_n\rangle = \langle \prod_{i=1}^n (x-\gamma_i)\rangle$ Hence I is principal, and $C^\omega[0,1]$ is a PID. □

The way one shows the existence of non-maximal prime ideals in the ring $C[0,1]$ is as follows: Any maximal element of the set of ideals disjoint from the multiplicatively closed set of non-zero polynomials partially ordered by inclusion can be checked to be a non-maximal prime ideal of $C[0,1]$. Let P be one such maximal element. Further, any prime ideal of $C[0,1]$ is contained in a 'unique' maximal ideal of $C[0,1]$ [Pan16]. (The above outline of the proof through Zorn's lemma is perhaps the only way to show the existence of non-maximal prime ideals in $C[0,1]$.)

Corollary. *The non maximal prime ideal P as mentioned above contains no nonzero analytic functions.*

Proof. Consider the contraction of P in the ring $C^\omega[0,1]$. It has to be a prime ideal of $C^\omega[0,1]$. Let M_γ be the unique maximal ideal of $C[0,1]$ such that $P \subset M_\gamma$. Then, the contraction of P in the ring $C^\omega[0,1]$ is either 0 or $M^\omega_\gamma = \langle x-\gamma\rangle$. But $x-\gamma$ is not in P, therefore P contracts to the zero ideal in $C^\omega[0,1]$. □

The fact that $C^\omega[0,1]$ is a PID while $C[0,1]$ has infinitely many uncountably generated ideals gives us an algebraic evidence to our intuition that real analytic functions are sparse as compared to real continuous functions (on a compact set).

References

[1] B. Sury, *Uncountably Generated Ideals of Functions*, The College Mathematics Journal,vol. 42, 404–406, 2011,Mathematical Association of America

[2] Vaibhav Pandey *On the prime ideals of C [0, 1]*, Resonance,Vol 21, No. 5, 439–446, 2016, Springer